

**1050**

# HOW TO IMPROVE COCHLEAR IMPLANT IN ADULT

## Poor performances in adult CI patients and its remediation

■ B. FRAYSSE



XXXVI Congreso

LIMA  
November 14-17, 2018

# GOAL OF THE STUDY



① To propose a predictive model during counselling based on patient related factors and electrode insertion

②


To compare early auditory outcomes to the predictive model and propose personalized remediation

# POPULATION

**N : 118**

## Inclusion

- All adults with unilateral CI and profound HL at least one year follow-up and receiving the same aural rehabilitation program



Cone Beam /  
CT-Scan

## Study design

- The percentage of variance (22) expresses the impact of each factors



EARLY SENTENCE RECOGNITION IN ADULT COCHLEAR IMPLANT USERS  
Chris James, Chadlia Karoui, Mathieu Marx, Marie-Laurence Laborde, Charles-Edouard Molinier, Benoit Lepage, Olivier Deguine, Bernard Escudé, Bernard Fraysse


*Accepted for publication*

# GOAL OF THE STUDY




- ① To analyze CI auditory outcomes as a function of delay post activation and the various factors underlying the results
- ② To design a predictive model during counselling based on patient related factors and electrode insertion
- ③ To compare early auditory outcomes to the predictive mode and propose remediation

# SENTENCES IN QUIET (1 DAY / 1 MONTH)




# INITIAL SCORE / 12 MONTHS




# OVERALL RESULTS

N = 118

Quiet




Noise 10 dB SNR





# The development of speech understanding with CI does not follow a linear function with time



# FACTORS TO BE CONSIDERED



## ■ Biographic and audiologic factors

- Age at implantation
- Etiology
- Duration of hearing loss



*Patient related*

## ■ Anatomical and surgical factors

- Insertion depth of apical electrodes
- Scala location



*Insertion technique*

## ■ Linguistic and neurocognitive skills



*Personalized auditory rehabilitation*



# BIOGRAPHIC FACTORS



- Age at implantation : **NS**
- Duration of deafness : 9 to 12% total variance  
(*0.46 pts per year of profound HL*)
- Etiologies : 20 to 30% total variance  
(*Chronic otitis, Meniere diseases*)


# BIOGRAPHIC FACTORS DATA

Blamey, P. J., Artières, F., Başkent, D., Bergeron, F., Beynon, A., Burke, E., ... Lazard, D. S. (2012). Factors Affecting Auditory Performance of Postlinguistically Deaf Adults Using Cochlear Implants: An Update with 2251 Patients. *Audiology & Neuro-Otology*, 2013;18(1): 36–47.



# ANATOMICAL AND SURGICAL FACTORS BASED ON POSTOPERATIVE CONE BEAM

## 1 Insertion depth




## 2 Scalar location




*In vivo* CT images: Professor Tobias Struffert, University of Erlangen.

# THE EFFECT OF INSERTION DEPTH ON AUDITORY OUTCOMES






# FREQUENCY ALLOCATION INSERTION DEPTH

- Moderate shifts may be easily accommodated but larger shifts  $> 1,5$  octave may affect auditory performance and the adaptation process take more time (e.g. *Li et al., 2009*)



# FREQUENCY ALLOCATION AND TONOTOPIC ORGANIZATION

- Size of the cochlea
- Type of electrode array
- Spacing between electrodes






# SCALAR LOCATION

	Scala tympani	Scala vestibuli or Dislocation
■ Type of electrode		
• Straight ( <i>N</i> : 43*)	38 (88%)	5 (12%)
• Perimodiolar ( <i>N</i> : 53**)	33 (62%)	20 (38%)
■ Depth of insertion	432°	403°



$p < 0.01$   
NS



# SCALAR DISLOCATION IMPACT AUDITORY OUTCOMES

2

- In our study the scala dislocation reduced scores by 12 - 25 pts at one year ( $p < 0.01$ ),  $r^2 = 14\%$



So why use a perimodiolar electrode ?

# VARIANCE OF PERIPHERAL FACTORS ON OUTCOMES

	In noise	In quiet
Etiology	0.34***	0.25**
Duration of deafness per year	0.06* per year	0.08* per year
Insertion length per degree	0.09***	0.08**
Proportion of electrodes in the scala tympani	0.14**	0.13**
<b>Total impact of peripheral factors</b>	<b>41%</b>	<b>49%</b>


# GOAL OF THE STUDY



- ① To analyze CI auditory outcomes as a function of delay post activation and the various factors underlying the results
- ② To design a predictive model during counselling based on patient related factors and electrode insertion
- ③ To compare early auditory outcomes to the predictive mode and propose remediation

# PREDICTIVE MODEL OF AUDITORY PERFORMANCE

- Based on our biographic data, we may develop a mathematical model during counselling based on biographic factors
  - 90 – 0.5/yr HL – (X étiologies)





- But also by an improvement of electrode insertion and reprogramming

# HOW CAN WE OPTIMIZE THE ELECTRODE INSERTION ?

Pre op.

- By a better surgical planning based on radiological data and electrode type




# HOW CAN WE OPTIMIZE THE ELECTRODE INSERTION ?



Pre op.


- By reprogramming the electrodes based on post operative insertion angle



El	Angle	SG Freq	Filtre Freq	Shift Oct
1	535	335,4	149	1,17
2	460	498,3	261	0,93
3	390	724,3	408	0,83
4	325	1047,5	601	0,80
5	270	1471,4	854	0,78
6	225	1993,7	1191	0,74
7	185	2674,6	1638	0,71
8	145	3680,3	2233	0,72
9	110	4974,1	3028	0,72
10	70	7192,2	4090	0,81
11	35	10159,1	5510	0,88
12	10	13327,4	7175	0,89


Decalage  
Moyen 0,83

# NEUROCOGNITIVE AND LINGUISTIC SKILLS



- In our study, 50 % of the variance at 1 month cannot be explained by auditory peripheral factors
- Speech discrimination in degraded condition (CI or HA to some limit) may be compensated by neurocognitive and linguistic skills
- The evolution of crossmodal plasticity is one of the underlying processes of compensatory mechanisms







## Evolution of Crossmodal Reorganization of the Voice Area in Cochlear-Implanted Deaf Patients

Julien Rouger,<sup>1</sup> Sébastien Lagleyre,<sup>2</sup> Jean-François Démonet,<sup>3</sup>  
Bernard Fraysse,<sup>2</sup> Olivier Deguine,<sup>1,2</sup> and Pascal Barone<sup>1\*</sup>


- We studied the dynamics of reversed crossmodal plasticity by TEP Brain imaging during auditory speech tracking



- Auditory stimulation determined a reactivation of auditory cortical areas but also a crossmodal reorganization of the cortical visual network



- Clear correlation between individual visual and auditory brain activity and auditory outcomes



# THE IMPORTANCE OF AUDITORY AND COGNITIVE REHABILITATION STRATEGIES

*Laryngoscope Investigative Otolaryngology*  
© 2016 The Authors Laryngoscope Investigative Otolaryngology  
published by Wiley Periodicals, Inc. on behalf of The Triological Society

---

## Postoperative Rehabilitation Strategies Used by Adults With Cochlear Implants: A Pilot Study

---

Michael S. Harris, MD; Natalie R. Capretta, BS; Shirley C. Henning, MS, CCC-SLP; Laura Feeney, AuD;  
Mark A. Pitt, PhD; Aaron C. Moberly, MD

- This article confirm the interest to develop specific rehabilitation strategies according to early outcomes in auditory and audiovisual conditions

# ON WHICH BASIS DEVELOP SPECIFIC REHABILITATION STRATEGIES?



- Optimization of the rehabilitation must take into account the level of :

① Phonemic sensitivity and lexical knowledge

② Cognitive factors

→ Speed of processing

→ Working memory and attention


→ Executive function

} Depend

One month results → Normal performer → Conventional rehabilitation



Poor performer



Auditory evaluation on objective method

- Frequency allocation
- Scalar location
- Speed of stimulation refractory period
- Channel selectivity spread of excitation

} X-Ray


Working memory and phonological sensitivity evaluation



Personalized auditory rehabilitation




« Bottom up » approach



« Top down » approach



# PERSONALIZED REHABILITATION PROGRAM



# TIMING OF INTERVENTION

Predictive Model Counselling patient	Electrode insertion Surgical planning	Optimized Fitting	Sentence recognition score Compare to predictive model	SNR50 as expected ?
Hearing story Etiology	Size of cochlea Type of electrode	X-Ray electrophysiological Frequency allocation	<ul style="list-style-type: none"> <li>❶ Data log</li> <li>❷ Lexical knowledge</li> <li>❸ Spread of excitation Recovery period</li> </ul>	Data log Loudness growth
		→ Evaluate 2 programs	<b>1 MONTH</b>	<b>6 MONTHS</b>
		<b>ACTIVATION</b>		
	<b>SURGERY</b>			
<b>PRE OP</b>			Cognitive or auditory } Training	→ MAP rehabilitation refinement

# CONCLUSION



- Counseling patients with realistic expectations and take into account the patient's goals (GAS)
- Electrode insertion should avoid dislocation and be adapted to the tonotopic organization
- Develop personalized rehabilitation programs and material based on early outcomes and targeted on specific weaknesses





***Thank you for your attention***